



TITLE:

HARDY CLASS OF FUNCTIONS DEFINED BY SALAGEAN OPERATOR

AUTHOR(S):

NIWA, NORIO; JIMBO, TOSHIYA; OWA, SHIGEYOSHI

CITATION:

NIWA, NORIO ...[et al]. HARDY CLASS OF FUNCTIONS DEFINED BY SALAGEAN OPERATOR.
数理解析研究所講究録 1996, 963: 67-73

ISSUE DATE:

1996-08

URL:

<http://hdl.handle.net/2433/60559>

RIGHT:

HARDY CLASS OF FUNCTIONS DEFINED BY SALAGEAN OPERATOR

NORIO NIWA, TOSHIYA JIMBO AND SHIGEYOSHI OWA

丹羽典朗 (奈良教育大) 神保敏弥 (奈良教育大) 尾和重義 (近畿大・理工)

ABSTRACT. The object of the present paper is to derive some properties for Hardy class of analytic functions defined by Salagean operator.

1. INTRODUCTION

Let A be the class of functions $f(z)$ of the form

$$(1.1) \quad f(z) = z + \sum_{k=2}^{\infty} a_k z^k$$

that are analytic in the open unit disk $U = \{z : |z| < 1\}$.

For $f(z) \in A$, the Salagean operator D^n (cf. [6]) is defined by

$$(1.2) \quad D^0 f(z) = f(z),$$

$$(1.3) \quad D^1 f(z) = Df(z) = zf'(z),$$

$$(1.4) \quad D^n f(z) = D(D^{n-1}f(z)) \quad (n \in \mathbb{N} = \{1, 2, 3, \dots\}).$$

A function $f(z)$ belonging to A is said to be starlike of order α if it satisfies

$$(1.5) \quad \operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} > \alpha \quad (z \in U)$$

for some α ($0 \leq \alpha < 1$). We denote by $S^*(\alpha)$ the subclass of A consisting of functions which are starlike of order α in U .

A function $f(z) \in A$ is said to be convex of order α if it satisfies

$$(1.6) \quad \operatorname{Re} \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \alpha \quad (z \in U)$$

for some α ($0 \leq \alpha < 1$). Also we denote by $K(\alpha)$ the subclass of A consisting of all such functions. Note that $f(z) \in K(\alpha)$ if and only if $zf'(z) \in S^*(\alpha)$ for $0 \leq \alpha < 1$.

Let H^p ($0 < p \leq \infty$) be the class of all analytic functions in U such that

$$(1.7) \quad \|f\|_p = \lim_{r \rightarrow 1^-} \{M_p(r, f)\} < \infty,$$

where

$$(1.8) \quad M_p(r, f) = \begin{cases} \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{\frac{1}{p}} & (0 < p < \infty) \\ \max_{|z| \leq r} |f(z)| & (p = \infty) \end{cases} \quad (\text{cf. [1]}).$$

AMS (1991) Subject Classification. Primary 30C45.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$

2. SOME LEMMAS

To discuss our problems for Hardy class H^p of functions, we need the following lemmas.

Lemma 1 ([7]). If $f(z) \in K(\alpha)$, then $f(z) \in S^*(\beta)$, where

$$(2.1) \quad \beta = \beta(\alpha) = \begin{cases} \frac{1-2\alpha}{2(2^{1-2\alpha}-1)} & (\alpha \neq \frac{1}{2}) \\ \frac{1}{2 \log 2} & (\alpha = \frac{1}{2}). \end{cases}$$

This result is sharp.

Lemma 2 ([2]). If $f(z) \in S^*(\alpha)$ and is not of the form

$$(2.2) \quad f(z) = \frac{z}{(1 - ze^{it})^{2(1-\alpha)}},$$

then there exists $\delta = \delta(f) > 0$ such that $\frac{f(z)}{z} \in H^{\delta + \frac{1}{2(1-\alpha)}}$.

Lemma 3 ([5]). If $p(z)$ is analytic in U with $p(0) = 1$ and

$$(2.3) \quad \operatorname{Re}(p(z) + zp'(z)) > \frac{1 - 2 \log 2}{2(1 - \log 2)} \quad (z \in U),$$

then $\operatorname{Re}(p(z)) > 0$ ($z \in U$).

Remark. We see that

$$\frac{1 - 2 \log 2}{2(1 - \log 2)} = -0.629 \dots$$

Lemma 4 ([1]). Every analytic function $p(z)$ with positive real part in U is in the class H^p for all $0 < p < 1$.

Lemma 5 ([4]). If $f(z) \in A$ satisfies $z^r f(z) \in H^p$ ($0 < p < \infty$) for a real r , then $f(z) \in H^p$ ($0 < p < \infty$).

Lemma 6 ([1]). If $f'(z) \in H^p$ for some p ($0 < p < 1$), then $f(z) \in H^q$ ($q = p/(1-p)$).

Lemma 7 ([3]). Let $w(z)$ be analytic in U with $w(0) = 0$. If $|w(z)|$ attains its maximum value on the circle $|z| = r$ ($0 \leq r < 1$) at a point z_0 , then we can write

$$z_0 w'(z_0) = k w(z_0),$$

where k is real and $k \geq 1$.

3. HARDY CLASS OF FUNCTIONS

Our first result for Hardy class is contained in

Theorem 1. *Let $f(z) \in A$ satisfy*

$$(3.1) \quad \operatorname{Re} \left\{ \frac{D^{n+1}f(z)}{D^n f(z)} \right\} > \alpha_0 \quad (z \in U)$$

for some α_0 ($0 \leq \alpha_0 < 1$), and let

$$(3.2) \quad \alpha_j = \begin{cases} \frac{1 - 2\alpha_{j-1}}{2(2^{1-2\alpha_{j-1}} - 1)} & (\alpha_{j-1} \neq \frac{1}{2}) \\ \frac{1}{2 \log 2} & (\alpha_{j-1} = \frac{1}{2}) \end{cases}$$

for $j = 1, 2, \dots, n$. If $D^{n-j}f(z)$ is not of the form

$$(3.3) \quad D^{n-j}f(z) = \frac{z}{(1 - ze^{it})^{2(1-\alpha_j)}},$$

then there exists $\delta > 0$ such that $D^{n-j}f(z) \in H^{\delta + \frac{1}{2(1-\alpha_j)}}$.

Proof. Note that

$$(3.4) \quad \begin{aligned} D^{n+1}f(z) &= D(D^n f(z)) \\ &= z(D^n f(z))' \\ &= z(D^{n-1}f(z))' + z^2(D^{n-1}f(z))'' \end{aligned}$$

and

$$(3.5) \quad D^n f(z) = z(D^{n-1}f(z))'.$$

This implies that

$$(3.6) \quad \operatorname{Re} \left\{ \frac{D^{n+1}f(z)}{D^n f(z)} \right\} = \operatorname{Re} \left\{ 1 + \frac{z(D^{n-1}f(z))''}{(D^{n-1}f(z))'} \right\} > \alpha_0,$$

so that, $D^{n-1}f(z) \in K(\alpha_0)$. Therefore, an application of Lemma 1 leads to

$$\begin{aligned} D^{n-1}f(z) \in K(\alpha_0) &\implies D^{n-1}f(z) \in S^*(\alpha_1) \\ &\iff D^{n-2}f(z) \in K(\alpha_1) \\ &\implies D^{n-2}f(z) \in S^*(\alpha_2) \\ &\dots \\ &\iff D^{n-j}f(z) \in K(\alpha_{j-1}) \\ &\implies D^{n-j}f(z) \in S^*(\alpha_j). \end{aligned}$$

Further, by using Lemma 2 and Lemma 5, we know that there exists $\delta > 0$ such that $D^{n-j}f(z) \in H^{\delta + \frac{1}{2(1-\alpha_j)}}$. ■

Taking $j = n$ in Theorem 1, we have

Corollary 1. Let $f(z) \in A$ satisfy (3.1) for some α_0 ($0 \leq \alpha_0 < 1$), and let

$$\alpha_n = \begin{cases} \frac{1 - 2\alpha_{n-1}}{2(2^{1-2\alpha_{n-1}} - 1)} & (\alpha_{n-1} \neq \frac{1}{2}) \\ \frac{1}{2 \log 2} & (\alpha_{n-1} = \frac{1}{2}). \end{cases}$$

If $f(z)$ is not of the form (3.3), then there exists $\delta > 0$ such that $f(z) \in H^{\delta + \frac{1}{2(1-\alpha_n)}}$.

Next, we derive

Theorem 2. Let $f(z) \in A$ satisfy

$$(3.7) \quad \operatorname{Re} \left\{ \frac{D^{n+1}f(z)}{z} \right\} > \frac{1 - 2 \log 2}{2(1 - \log 2)} \quad (z \in U).$$

Then there exists p_j ($j = 1, 2, \dots, n+1$) such that $D^{n-j+1}f(z) \in H^{p_j}$, where

$$(3.8) \quad p_k < \frac{1}{j - k + 1} \quad (k = 1, 2, \dots, j).$$

Proof. Define the function $p(z)$ by

$$(3.9) \quad p(z) = \frac{D^n f(z)}{z}.$$

Then $p(z)$ is analytic in U and $p(0) = 1$. Since

$$(3.10) \quad \operatorname{Re} \left\{ \frac{D^{n+1}f(z)}{z} \right\} = \operatorname{Re}(p(z) + zp'(z)) > \frac{1 - 2 \log 2}{2(1 - \log 2)},$$

Lemma 3 gives that

$$(3.11) \quad \operatorname{Re}(p(z)) = \operatorname{Re} \left\{ \frac{D^n f(z)}{z} \right\} > 0 \quad (z \in U).$$

Noting that

$$\frac{D^n f(z)}{z} = (D^{n-1}f(z))',$$

an application of Lemma 4 implies that $(D^{n-1}f(z))' \in H^{p_1}$, so by Lemma 6,

$$D^{n-1}f(z) \in H^{p_2} \quad (p_2 = \frac{p_1}{1 - p_1}).$$

Further, since $D^{n-1}f(z) = z(D^{n-2}f(z))'$, using Lemma 5, we obtain $(D^{n-2}f(z))' \in H^{p_2}$. Taking this process again and again, we conclude that $D^{n-j+2}f(z) \in H^{p_{j-1}}$ and $0 < p_{j-1} < 1/2$. Thus, finally we have $D^{n-j+1}f(z) \in H^{p_j}$ ($0 < p_j < 1$). This completes the proof of Theorem 2. ■

Letting $j = n+1$ in Theorem 2, we have

Corollary 2. Let $f(z) \in A$ satisfy (3.7). Then there exists p_{n+1} such that $f(z) \in H^{p_{n+1}}$, where

$$p_k < \frac{1}{n - k + 2} \quad (k = 1, 2, \dots, n+1).$$

4. HARDY CLASS OF BOUNDED FUNCTIONS

Next our theorem for Hardy class of bounded functions is contained in

Theorem 3. Let $f(z) \in A$ satisfy

$$(4.1) \quad \left| \frac{D^{n+2}f(z)}{D^{n+1}f(z)} - 1 \right| < \frac{5\alpha_0 - 2\alpha_0^2 - 1}{2\alpha_0} \quad (z \in U)$$

for some α_0 ($1/3 \leq \alpha_0 \leq 1/2$), or

$$(4.2) \quad \left| \frac{D^{n+2}f(z)}{D^{n+1}f(z)} - 1 \right| < \frac{\alpha_0 - 2\alpha_0^2 + 1}{2\alpha_0} \quad (z \in U)$$

for some α_0 ($1/2 \leq \alpha_0 < 1$). If $D^{n-j}f(z)$ is not of the form (3.3), then there exists $\delta > 0$ such that $D^{n-j}f(z) \in H^{\delta + \frac{1}{2(1-\alpha_j)}}$ ($j = 1, 2, \dots, n$), where α_j is given by (3.2).

Proof. Define the function $w(z)$ by

$$(4.3) \quad \frac{D^{n+1}f(z)}{D^n f(z)} = \frac{1 + (1 - 2\alpha_0)w(z)}{1 - w(z)} \quad (w(z) \neq 1).$$

Then $w(z)$ is analytic in U and $w(0) = 0$. It follows from (4.3) that

$$(4.4) \quad \begin{aligned} & \frac{D^{n+2}f(z)}{D^{n+1}f(z)} - 1 \\ &= \left(\frac{w(z)}{1 - w(z)} \right) \left(2(1 - \alpha_0) + \frac{zw'(z)}{w(z)} + \frac{(1 - 2\alpha_0)(1 - w(z))}{1 + (1 - 2\alpha_0)w(z)} \left(\frac{zw'(z)}{w(z)} \right) \right). \end{aligned}$$

Suppose that there exists a point $z_0 \in U$ such that

$$\max_{|z| \leq |z_0|} |w(z)| = |w(z_0)| = 1 \quad (w(z_0) \neq 1).$$

Then Lemma 7 leads us to $w(z_0) = e^{i\theta}$ and

$$z_0 w'(z_0) = k w(z_0) \quad (k \geq 1).$$

Therefore, we have

$$(4.5) \quad \begin{aligned} & \left| \frac{D^{n+2}f(z_0)}{D^{n+1}f(z_0)} - 1 \right| \\ &= \left| \frac{w(z_0)}{1 - w(z_0)} \right| \left| 2(1 - \alpha_0) + \frac{zw'(z_0)}{w(z_0)} + \frac{(1 - 2\alpha_0)(1 - w(z_0))}{1 + (1 - 2\alpha_0)w(z_0)} \left(\frac{zw'(z_0)}{w(z_0)} \right) \right| \\ &= \left| \frac{e^{i\theta}}{1 - e^{i\theta}} \right| \left| 2(1 - \alpha_0) + k + k \frac{(1 - 2\alpha_0)(1 - e^{i\theta})}{1 + (1 - 2\alpha_0)e^{i\theta}} \right| \\ &\geq \frac{2(1 - \alpha_0) + k}{|1 - e^{i\theta}|} - \frac{k|1 - 2\alpha_0|}{|1 + (1 - 2\alpha_0)e^{i\theta}|} \\ &\geq \frac{2(1 - \alpha_0) + k}{2} - \frac{k|1 - 2\alpha_0|}{2\alpha_0}. \end{aligned}$$

For $1/3 \leq \alpha_0 \leq 1/2$, we have

$$(4.6) \quad \left| \frac{D^{n+2}f(z_0)}{D^{n+1}f(z_0)} - 1 \right| \geq \frac{5\alpha_0 - 2\alpha_0^2 - 1}{2\alpha_0}$$

and for $1/2 \leq \alpha_0 < 1$, we have

$$(4.7) \quad \left| \frac{D^{n+2}f(z_0)}{D^{n+1}f(z_0)} - 1 \right| \geq \frac{\alpha_0 - 2\alpha_0^2 + 1}{2\alpha_0}.$$

Since the above contradicts our conditions (4.1) and (4.2) of the theorem, we conclude that $|w(z)| < 1$ for all $z \in U$. This implies that

$$(4.8) \quad \operatorname{Re} \left\{ \frac{D^{n+1}f(z)}{D^n f(z)} \right\} > \alpha_0 \quad (z \in U).$$

Noting that (4.8) is equivalent to $D^n f(z) \in S^*(\alpha_0)$. Using the same manner in the proof of Theorem 1, we conclude that $D^{n-j}f(z) \in S^*(\alpha_j)$. Thus, applying Lemma 2 and Lemma 5, we can prove Theorem 3. ■

If we put $j = n$ in Theorem 3, then we have

Corollary 3. *Let $f(z) \in A$ satisfy the condition (4.1) for some α_0 ($1/3 \leq \alpha_0 \leq 1/2$) or (4.2) for some α_0 ($1/2 \leq \alpha_0 < 1$). If $f(z)$ is not of the form (3.3), then there exists $\delta > 0$ such that $f(z) \in H^{\delta + \frac{1}{2(1-\alpha_n)}}$, where α_n is given by (3.2).*

ACKNOWLEDGMENTS

This work of authors was supported, in part, by the Japanese Ministry of Education, Science and Culture under Grant-in-Aid for General Scientific Research.

REFERENCES

1. P. L. Duren, *Theory of H^p Spaces*, A series of Monographs and Textbooks in Pure and Applied Mathematics, vol. 38, Academic Press, New York and London, 1970.
2. P. J. Eenigenburg and F. R. Keogh, *The Hardy class of some univalent functions and their derivatives*, Michigan Math. J. 17 (1970), 335–346.
3. I. S. Jack, *Functions starlike and convex of order α* , J. London Math. Soc. 3 (1971), 469–474.
4. Y. C. Kim, K. S. Lee and H. M. Srivastava, *Certain classes of integral operators associated with the Hardy space of analytic functions*, Complex Variables 20 (1992), 1–12.
5. M. Nunokawa, *On starlikeness of Libera transformation*, Complex Variables 17 (1991), 79–83.
6. G. S. Salagean, *Subclasses of univalent functions*, Lecture Notes in Math. 1013, (C. A. Cazacu, N. Boboc, M. Jurchescu and I. Susiu Ed.), Springer-Verlag, Berlin, Heidelberg, New York and Tokyo, 1983, pp. 362–372.

7. D. R. Wilken and J. Feng, *A remark on convex and starlike functions*, J. London Math. Soc. **21** (1980), 287–290.

N. Niwa and T. Jimbo : Department of Mathematics, Nara University of Education, Takabatake, Nara 630, Japan.

S. Owa : Department of Mathematics, Kinki University, Higashi-Osaka, Osaka 577, Japan.